

On the existence of a rainbow 1-factor in proper coloring of $K_{rn}^{(r)}$ *

Xueliang Li¹ and Zhixia Xu^{1,2,†}

¹Center for Combinatorics and LPMC-TJKLC,
Nankai University, Tianjin 300071, P.R. China

²College of Mathematics and System Sciences,
Xinjiang University, Urumuqi, 830046, P.R. China

Abstract

El-Zanati et al proved that for any 1-factorization $\mathcal{F}$ of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$ with $r \geq 2$ and $n \geq 3$, there is a rainbow 1-factor. We generalize their result and show that in any proper coloring of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$ with $r \geq 2$ and $n \geq 3$, there is a rainbow 1-factor.

Keywords: edge-colored graph, rainbow 1-factor, rainbow matching

AMS Subject Classification 2000: 05C15, 05C35, 05C55, 05C70.

1 Introduction

A hypergraph $\mathcal{G} = (\mathcal{V}, \mathcal{E})$ consists of a finite set $\mathcal{V}$ of vertices and a set $\mathcal{E}$ of subsets of $\mathcal{V}$ called edges. An edge subset $\mathcal{E}'$ of disjoint edges of $\mathcal{E}$ is called *independent*. A proper coloring of $\mathcal{E}$ is a partition of $\mathcal{E}$ into independent sets with each partition set is given a color, say $1, 2, \dots, l$. For a given coloring of $\mathcal{G}$, a subhypergraph $\mathcal{G}'$ is called *rainbow* if each edge of $\mathcal{G}'$ has distinct color. A 1-factor of a hypergraph $(\mathcal{V}, \mathcal{E})$ is an independent edge set which partition

*Supported by NSFC, PCSIRT and the “973” program. Email: lxl@nankai.edu.cn (Xueliang Li); irisxuzx@gmail.com (Zhixia Xu). † corresponding author

$\mathcal{V}$. A 1-factorization of $(\mathcal{V}, \mathcal{E})$ is a partition of $\mathcal{E}$ into 1-factors. For positive integers $r \geq 2$ and n , the complete r -uniform hypergraph on n vertices is the hypergraph $K_n^{(r)}$, with a vertex set $\mathcal{V}$ of order n and an edge set $\mathcal{E}$ consisting of all r -subsets of $\mathcal{V}$. Note that $K_n^{(2)}$ is K_n , the simple complete graph of order n . In order for $K_n^{(r)}$ to contain a 1-factor, it is clearly necessary that r divides n . In 1973, Baranyai [1] showed that $K_{nr}^{(r)}$ has a 1-factorization. Given a 1-factorization $\mathcal{F}$ of $K_{nr}^{(r)}$, $n \geq 3$, Woolbright in 1978 showed that there exists a 1-factor in $K_{nr}^{(r)}$ whose edges belong to at least $n - 1$ different 1-factors of $\mathcal{F}$. In 1998, Woolbright and Fu [4] proved that for any 1-factorization of K_{2n} there is a rainbow 1-factor.

In [2], El-Zanati et al proved that for any 1-factorization $\mathcal{F}$ of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$ with $r \geq 2$ and $n \geq 3$, there is a rainbow 1-factor. It is clear that a 1-factorization is a very special case of proper colorings. In the present paper, we want to use a weaker condition, the proper coloring condition, to replace their stronger condition, the 1-factorization condition, and to generalize their result as follows: for any proper coloring of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$ with $r \geq 2$ and $n \geq 3$, there is a rainbow 1-factor. To show the result, we divide the proof into three cases: $r = 2$ and $n \geq 3$, $r > 2$ and $n = 3$, $r > 2$ and $n > 3$. Notice that the proof of Theorem 3 for the case $r > 2$ in [2] can be used directly to show that for $r > 2, n > 3$ there is a rainbow 1-factor in any proper coloring of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$. For the case $r > 2, n = 3$, we give a prove in Theorem 2.3. The substantial part of our proof is to show the result for the case $r = 2$ and $n \geq 3$, which will be given in Theorems 2.1 and 2.2. As a result we have

Theorem 1.1. *For any proper coloring of the complete uniform hypergraph $\mathcal{G} = K_{rn}^{(r)}$ with $r \geq 2$ and $n \geq 3$, there is a rainbow 1-factor.*

2 Existence of a rainbow 1-factor in proper coloring of K_{2n} and $K_{3r}^{(r)}$

Let $G = (V, E)$ be a graph and C be a proper coloring of G . A *rainbow matching* of G is a matching whose edges have pairwise different colors. For $e \in E$, let $C(e)$ denote the color of e . For $v \in V$, let $C(v) = \{C(e) | e \text{ is incident with } v\}$. For any subset E' of E , let $C(E') = \{C(e) | e \in E'\}$ and $F(E') = C(E) - C(E')$.

For any $V' \subseteq V$, let $G[V']$ denote the subgraph induced by V' .

Lemma 2.1. *For any proper coloring of K_{2n} , there is a rainbow perfect matching when $n = 3$ or $n = 4$.*

Proof. For $n = 3$, let the vertices of K_6 be $v_1, v_2, \dots, v_6$ and C be a proper coloring of K_6 , let $1, 2, \dots, l$ be the colors used, we will show that there is a rainbow $3K_2$ in C . Suppose 1 is the color that appears least times in C on $E(K_6)$. If 1 appears on three edges, then C is a 1-factorization of K_6 and by the proof in [1], there is a rainbow 1-factor in C . If 1 appears on two edges, say $C(x_1x_2) = C(x_3x_4) = 1$, assume that $C(x_5x_1) = 2, C(x_5x_2) = 3, C(x_5x_3) = 4, C(x_5x_4) = 5, C(x_5x_6) = 6$. Since $\{x_5x_1, x_6x_2, x_3x_4\}$ is independent, to avoid the existence of rainbow perfect matching, it must be that $C(x_6x_2) = 2$. Similarly, $\{x_5x_2, x_6x_1, x_3x_4\}$ is independent and $C(x_6x_1) = 3$; $\{x_1x_2, x_5x_3, x_6x_4\}$ is independent and $C(x_6x_4) = 4$; $\{x_1x_2, x_5x_4, x_6x_3\}$ is independent and $C(x_6x_3) = 5$. Now both $\{x_5x_2, x_6x_3, x_1x_4\}$ and $\{x_5x_3, x_6x_2, x_1x_4\}$ are independent, whatever color the edge x_1x_4 receives, we will have a rainbow perfect matching. If 1 appears only once in C and $c(x_1x_2) = 1$, to avoid the existence of rainbow perfect matching, there is no rainbow $2K_2$ in the subgraph induced by $\{v_2, v_3, v_4, v_5\}$. The only such coloring is $c(x_3x_4) = c(x_5x_6) = 2, c(x_3x_5) = c(x_4x_6) = 3, c(x_3x_6) = c(x_5x_4) = 4$. Assume $c(x_3x_1) = 5$. Since both $\{x_1x_3, x_5x_6, x_2x_4\}$ and $\{x_1x_3, x_4x_6, x_2x_5\}$ are independent, we have $C(x_2x_4) = 5$ and $C(x_2x_5) = 5$, a contradiction. So in any proper coloring of K_6 , there is a rainbow perfect matching.

For $n = 4$, let the vertices of K_8 be $v_1, v_2, \dots, v_8$ and C be a proper coloring of K_8 , we will show that there is a rainbow $4K_2$ in C . Starting with any triangle, it is possible to find in C of $G = K_8$ at least two rainbow K_4 , say $G[\{v_1, v_2, v_3, v_4\}]$ and $G[\{v_1, v_2, v_3, v_5\}]$ are both rainbow. If there is at least one rainbow $2K_2$ in $G[\{v_5, v_6, v_7, v_8\}]$, since $\{v_1v_2, v_3v_4\}, \{v_1v_3, v_2v_4\}$ and $\{v_1v_4, v_2v_3\}$ are all independent and each edge has a distinct color, we can find a rainbow $4K_4$ in C . Similarly, there is no rainbow $2K_2$ in $G[\{v_4, v_6, v_7, v_8\}]$. But it is impossible that both $G[\{v_5, v_6, v_7, v_8\}]$ and $G[\{v_4, v_6, v_7, v_8\}]$ have no rainbow $2K_2$, and the proof is complete. $\square$

Theorem 2.2. *For $n \geq 3$, any proper coloring of K_{2n} contains a rainbow perfect matching.*

Proof. By Lemma 1, we can assume $n \geq 5$. Let C be any proper coloring of K_{2n} with the colors named $1, 2, \dots, l$, $l \geq 2n - 1$. Let $\mathcal{M}$ be any maximal rainbow matching with $|\mathcal{M}| = k$. Suppose $k < n$, we will show that there must be a rainbow matching with $k + 1$ edges. Recall that $C(\mathcal{M})$ denotes the set of colors of $\mathcal{M}$ and $F(\mathcal{M})$ denotes the complementary set of colors. Let s, t be two unmatched vertices. We may assume that $C(\mathcal{M}) = \{1, 2, \dots, k\}$ and $C(st) = 1$. Note that by maximality of $\mathcal{M}$, any edge incident with s whose color is in $F(\mathcal{M})$ must be incident with an edge of $\mathcal{M}$.

Let $C(s) = C_1 \cup C_2$ with $C_1 \subseteq C(\mathcal{M})$, $C_2 \subseteq F(\mathcal{M})$, $|C_1| = p \leq k$, $|C_2| = 2n - 1 - p$. Consider all the (s, t) -paths of length three, whose first edge is colored with a color α in $F(\mathcal{M})$, and the second edge is in $\mathcal{M}$; we call them the candidate 3-paths relative to $\mathcal{M}$. We can assume that each of these paths has its third edge colored with a color either in $C(\mathcal{M}) - \{1\}$, or the color α again, for otherwise we could augment $\mathcal{M}$ to $k + 1$ edges simply by deleting the second edge of the path from it and adding the first and third edges. There are $2n - 1 - p$ of these candidate paths and only $k - 1$ colors in $C(\mathcal{M}) - \{1\}$. So it follows that at least $2n - p - k$ of these paths have the first and third edges colored with the same color in $F(\mathcal{M})$; we call such paths $\mathcal{M}$ -symmetric (s, t) -paths.

Let $C(t) = C'_1 \cup C'_2$ with $C'_1 \subseteq C(\mathcal{M})$, $C'_2 \subseteq F(\mathcal{M})$, $|C'_1| = q \leq k$, $|C'_2| = 2n - 1 - q$. Consider the $2n - 1 - q$ edges incident with t whose colors are in $F(\mathcal{M})$. Each of these edges must be incident with an edge of $\mathcal{M}$, by the maximality of $\mathcal{M}$; at most 2 of them, say the ones colored $k + 1$ and $k + 2$, are incident with the edge of $\mathcal{M}$ colored 1. Now let $L = C(t) \setminus (C(\mathcal{M}) \cup \{k + 1, k + 2\})$ and $|L| = 2n - q - 3$.

For each color $i \in L$, we define a slight variation of the $(\mathcal{M}, st)$ pair. If the edge of color i incident with vertex t is $e_t = \{t, z_i\}$ of $\mathcal{M}$, we let the corresponding matching be $\mathcal{M}_i = (\mathcal{M} - \{e_i\}) \cup \{e_t\}$; now t_i is unmatched (in $\mathcal{M}_i$), and we let our starting/ending vertex pair be s, t_i , respectively. Note that $F(\mathcal{M}_i) = (F(\mathcal{M}) - i) \cup \{C(e_i)\}$. Also note that $C(e_i) \neq 1$, because i is neither $k + 1$ or $k + 2$.

As in the previous discussion, for each such i , there are $2n - 1 - p$ candidate 3-paths relative to $\mathcal{M}_i$ starting at s , ending at t_i , whose first edge is colored with a color in $F(\mathcal{M}_i)$, and whose second edge is in $\mathcal{M}_i$. Again, we assume that at least $2n - p - k$ of these paths are symmetric. Thus, listing the symmetric

paths for $i \in L$, we get a total of at least $(2n - q - 3)(2n - p - k)$ paths in the list of symmetric candidate paths. However, because in each of these symmetric paths either the middle edge is in $\mathcal{M}$, or the path has the form $sz_i tt_i$, and therefore has the same first and third edges as $sz_i t_i t$, each of the symmetric candidate paths is uniquely determined by its first edge. Moreover, the color α of the starting/ending edge in these symmetric paths cannot be $c(e_i) = c(\{z_i, t_i\})$ (the only possible such path has vertex sequence $stz_i t_i$, which is not symmetric because $c(e_i) \neq 1$), so α must be in C_2 . Therefore each of the possible starting color can only start one path in the list. It follows that $2n - p - 1 \geq (2n - q - 3)(2n - p - k)$. Let $x = 2n - p$, $y = 2n - q$, then $x - 1 \geq (y - 3)(x - k)$, and $x(y - 4) < k(y - 3)$. Since $q \leq k < n$, $y = 2n - q = n + n - q \geq n + n - k > n$, $y > 4$. Then we have $\frac{x}{k} < \frac{y-3}{y-4}$ and $x < k$, that is $p + k > 2n$, which is contrary to $p \leq k < n$.

We conclude that there must be a rainbow matching with $k + 1$ edges, and so the result follows. $\square$

For the case $n = 3$, it is easy to see that in any 1-factorization of $K_{3r}^{(r)}$ there is a rainbow 1-factor, and the proof was omitted in [2]. But in a proper coloring, the proof is not straightforward, and we prefer to give the details in the following.

Theorem 2.3. *For $r \geq 2$, any proper coloring of $K_{3r}^{(r)}$ contains a rainbow perfect matching.*

Proof. Let the vertex set of $K_{3r}^{(r)}$ be $\mathcal{V} = \{x_1, x_2, \dots, x_{3r}\}$ and C be a proper coloring of $K_{3r}^{(r)}$. Take any two independent edges having a same color 1, say $m_1 = \{x_1, x_2, \dots, x_r\}$ and $m_2 = \{x_{r+1}, x_{r+2}, \dots, x_{2r}\}$. Then for any edge $m^1 \subset \mathcal{V} - m_1$ other than m_2 and $m^{1*} = \mathcal{V} - (m_1 \cup m^1)$, m^1 and m^{1*} have the same color and there is no other edge in this color, otherwise $\{m_1, m^1, m^{1*}\}$ is a rainbow 1-factor. Similarly, for any edge $m^2 \subset \mathcal{V} - m_2$ other than m_1 and $m^{2*} = \mathcal{V} - (m_2 \cup m^2)$, m^2 and m^{2*} have the same color and there is no other edge in this color. Let $m^1 = \{x_{r+1}, x_{r+2}, \dots, x_{2r-1}, x_{2r+1}\}$, $m^2 = \{x_1, x_{2r+2}, x_{2r+3}, \dots, x_{3r}\}$, then $\{m^1, m^2, \{x_2, x_3, \dots, x_r, x_{2r}\}\}$ is a rainbow 1-factor. $\square$

References

- [1] Zs. Baranyai, On the factorization of the complete uniform hypergraph, In: Infinite and finite sets I Colloq Math Soc János Bolyai 10, North-Holland, Amsterdam, 1975, pp.91-108.
- [2] S.I. El-Zanati, M.J. Plantholt, P.A. Sissokho and L.E. Spence, On the existence of rainbow 1-factor in 1-factorizations of $K_{rn}^{(r)}$, J. Combin. Des. 2007
- [3] D.E. Woolbright, On the size of partial 1-factor of 1-factorization of the complete k -uniform hypergraph on kn vertices, Ars Combin. 6(1978), 185-192.
- [4] D.E. Woolbright and H.L. Fu, On the existence of rainbows in 1-factorizations of K_{2n} , J. Combin. Des. 6(1998), 1-20.